

gelatinoso cinctis, atro-fuscis, subopacis, 20–21 \approx 8; paraphysibus crassis, articulatis, ascos superantibus simplicibus vel sparse ramosis.

Hab. in stercore leporino prope Brunn Moraviæ, sociis *Sporormia intermedia*, *Sordaria macrospora* et *S. discospora*.

8. **Delitschia sordarioides** Speg. Fungi Arg. Pug. II, n. 73. — Pe. 2778

Peritheciis globoso-lenticularibus, $\frac{1}{3}$ – $\frac{1}{4}$ mill. diam., olivaceo-fuscis, glabris, matrici semiimmersis, ostiolo papillato coriaceo-atro ornatis, contextu membranaceo, parenchymatico, dilute fusco-atro; ascis longe cylindraceis, sursum truncato-rotundatis, basi in pedicellum longum, ac gracilem attenuatis, 150–170 \approx 10–14, octosporis, paraphysatis (?); sporidiis subdistichis, primo continuis, cylindraceis, hyalinis, dein ellipticis, fuligineo-sub-opacis, utrinque rotundato-truncatis, 14–15 \approx 7–8, excentrice (inferne) 1-septatis, ad septum non constrictis, loculis, jodi ope, nucleo-atro-fuligineo hyalino-ocellato ornatis, duas caudas cylindraceas hyalinas gerentibus, superiore sporidium æquantem; inferiorem pluries longiorem, 30–35 \approx 5.

Hab. in fimo vaccino secus Rio de la Plata cerca de la Recoleta, Argent. Amer. merid. — *Sordaria coprophilæ* affinis, sed septo recedens; an generi *Delitschiæ* certe adscribenda sit nonnihil dubius sum. Speg. l. c.

9. **Delitschia Winteri** Plowr. Brit. Fung. in Grev. II, p. 188, t. 25, 2779

fig. 1. — Peritheciis semiimmersis, v. immersis, ostiolo cylindraceo, crassiusculo, curvulo, tuberculato, apice truncato subinde curvulo; ascis cylindraceis, deorsum breve attenuato-stipitatis, 450 \approx 65–70, apice rotundatis, paraphysibus septulatis, hyalinis obvallatis, octosporis; sporidiis oblique monostichis, oblongis, constricto-1-sepaatis, 65–75 \approx 29–35, utrinque rotundatis, primo flavis, dein opace fuligineis, strato gelatinoso crasso obvolutis.

Hab. in fimo vaccino, ovino et cuniculorum, North Wootton Britanniae, Conegliano Ital. bor.

10. **Delitschia elephantina** Pass. Funghi Abiss. p. 190, t. V, fig. 14. 2780

— Peritheciis minutis conicis hyphis reptantibus brevibus paucis affixis; ascis subclavatis rectiusculis vel curvis vix pedicellatis 8-sporis; sporidiis obscure distichis fusiformibus septatis, apicibus acutis, ad septum vix constrictis, primo hyalinis, tandem fuscescentibus.

Hab. in stercore Elephantis, Keren Abyssiniæ (Beccari). — Asci (ex icone) 27 μ . circ. longi, sporidia 10 μ . circ. longa; sed dimensiones pro genere justo multo minores videntur.